



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47, 16/18, G01N 33/53, C12Q 1/68, C12N 15/11, 9/00		A2	(11) International Publication Number: WO 99/67386
			(43) International Publication Date: 29 December 1999 (29.12.99)
(21) International Application Number: PCT/US99/14036 (22) International Filing Date: 22 June 1999 (22.06.99) (30) Priority Data: 60/090,391 23 June 1998 (23.06.98) US 60/118,570 3 February 1999 (03.02.99) US 09/337,171 21 June 1999 (21.06.99) US (71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US). (72) Inventor: KENNEDY, Giulia, C.; 360 Castenada Avenue, San Francisco, CA 94116 (US). (74) Agents: POTTER, Jane, E., R.; Chiron Corporation, 4560 Horton Street, Emeryville, CA 94608 (US) et al.			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: DIFFERENTIALLY EXPRESSED GENES IN PANCREATIC CANCER AND DISPLASIA			
(57) Abstract			
<p>The present invention provides the art with the DNA coding sequences of polynucleotides that are up- or down-regulated in cancer and dysplasia. These polynucleotides and encoded proteins or polypeptides can be used in the diagnosis or identification of cancer and dysplasia. Inhibitors of the up-regulated polynucleotides and proteins can decrease the abnormality of cancer and dysplasia. Enhancing the expression of down-regulated polynucleotides or introducing down-regulated proteins to cells can decrease the growth and/or abnormal characteristics of cancer and dysplasia.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DIFFERENTIALLY EXPRESSED GENES IN PANCREATIC CANCER AND DISPLASIA

This application claims priority to provisional applications Serial No. 60/090,391, filed June 23, 1998, and Serial No. 60/118,570, filed February 3, 1999.

5 TECHNICAL AREA OF THE INVENTION

The invention relates to the area of diagnosis and treatment of pancreatic cancer and dysplasia. More specifically, it relates to polynucleotides which are differentially regulated in pancreatic cancer and dysplasia.

BACKGROUND OF THE INVENTION

10 Pancreatic cancer is the fifth leading cause of cancer death in the United States. According to the American Cancer Society, approximately 28,000 people will die of pancreatic cancer in the United States in 1998. A high risk of developing pancreatic cancer, without a corresponding increase in the risk of developing other cancers, may be passed along in some families. Pancreatic cancer is most likely caused
15 by an accumulation of mutations in specific cancer-causing genes. Pancreatic cancer is very aggressive and chemotherapeutic agents which may be active against other malignancies do not work effectively when used for pancreatic cancer.

The majority of cells in the pancreas are in the exocrine glands, which produce pancreatic enzymes, and in the ducts that carry the pancreatic enzymes to the
20 bile duct and to the small intestine. Cancers of the exocrine cells of the pancreas are usually adenocarcinomas. Pancreatic adenocarcinomas usually begin in the ducts of the pancreas, but may sometimes develop from the acinar cells. About 95% of cancers of the pancreas are adenocarcinomas. Less common cancers of the exocrine pancreas include adenosquamous carcinomas, squamous cell carcinomas, and giant cell
25 carcinomas.

Because pancreatic cancer is an aggressive cancer with very high mortality, there is a need in the art for genes that are up- or down-regulated in tumor progression. Such genes are useful for therapeutic purposes and for diagnosis of pancreatic as well as other cancers.

SUMMARY OF THE INVENTION

The invention provides isolated polynucleotides comprising coding regions or portions of genes whose expression is mis-regulated in cancer and dysplasia.

The invention also provides isolated proteins and protein fragments
5 whose expression is mis-regulated in cancer and dysplasia.

The invention further provides an antibody preparation which specifically binds to a polypeptide the expression of which is mis-regulated in cancer and dysplasia.

The invention provides a method for diagnosing cancer and dysplasia.

10 The invention still further provides therapeutic compositions useful for treating cancer and dysplasia.

These and other objects of the invention are provided by one or more of the embodiments described below. One embodiment of the invention provides isolated polynucleotides comprising at least twelve contiguous nucleotides selected from the
15 group of polynucleotide sequences as shown in SEQ ID NOs:1-15.

Another embodiment of the invention provides isolated polypeptides comprising at least six contiguous amino acids encoded by a polynucleotide selected from the group consisting of the polynucleotide sequences as shown in SEQ ID NOs:1-
20 15.

Even another embodiment of the invention provides an antibody preparation which specifically binds to a polypeptide comprising at least six contiguous amino acids encoded by a polynucleotide selected from the group of polynucleotide sequences as shown in SEQ ID NOs:1-15.

Yet another embodiment of the invention provides isolated nucleotide
25 probes consisting of a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:1-15.

Still another embodiment of the invention provides a method of diagnosing cancer. The amount of a polypeptide expressed from a polynucleotide having a sequence as shown in SEQ ID NO:12 in a test sample of tissue of a human
30 suspected of being cancerous is determined. The amount of said polypeptide is also determined in a human tissue which is normal. The determined amounts are then

compared. A test sample which contains less of the polypeptide than the normal tissue is identified as cancerous.

A further embodiment of the invention provides an additional method of diagnosing cancer. The amount of specific mRNA molecules in a test sample of tissue suspected of being cancerous and in a human tissue which is normal are determined. The mRNA molecules to be measured are complementary to the minus strand of a double-stranded polynucleotide sequence. The double-stranded polynucleotide sequence is shown in SEQ ID NO:12. The determined amounts of mRNA molecules are compared. A test sample of tissue which contains less of the mRNA molecules than the normal tissue is identified as cancerous.

Another embodiment of the invention provides a therapeutic composition useful for reducing the growth rate of cancer cells. The composition is comprised of a polynucleotide comprising all or a portion of a nucleotide sequence which is operably linked to a promoter sequence and a pharmaceutically acceptable carrier. The polynucleotide comprising all or a portion of a nucleotide sequence comprises at least 18 contiguous nucleotides. The nucleotide sequence is shown in SEQ ID NO:12.

Yet another embodiment of the invention provides a therapeutic composition useful for reducing the growth rate of cancer cells. The composition is comprised of a polypeptide comprising all or a portion of an amino acid sequence expressed from a polynucleotide sequence and a pharmaceutically acceptable carrier. The polynucleotide sequence is shown in SEQ ID NO:12.

Another embodiment of the invention provides a method of diagnosing dysplasia and cancer. The amount of a polypeptide expressed from a polynucleotide having at least one of a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:2, 5, and 15 in a test sample of tissue suspected of being dysplastic or cancerous is determined. The amount of the polypeptide is also determined in a human tissue which is normal. The determined amounts are compared. A test sample of human tissue which contains more of at least one polypeptide than the normal tissue is identified as being dysplastic or cancerous.

A further embodiment of the invention provides another method of diagnosing dysplasia. The amount of a polypeptide expressed from a polynucleotide having a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14 is determined in a test sample of tissue suspected of being dysplastic. The amount of said polypeptide is also determined in a human tissue which is normal. The two amounts are then compared. A test sample of human tissue which contains more of said polypeptide than the normal tissue is identified as being dysplastic.

Another embodiment of the invention provides an additional method of diagnosing cancer.

The amount of a polypeptide expressed from a polynucleotide having a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:2, 5, and 15, is determined in a test sample of tissue suspected of containing cancer, and in a human tissue which is normal. The amount of a polypeptide expressed from a polynucleotide having a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14 is also determined in the test sample, and in the normal tissue. The determined amounts of said polypeptides are then compared. A test sample of tissue which contains more of the polypeptide expressed from a polynucleotide having a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:2, 5, and 15, as compared to the normal tissue, and which contains substantially the same amount of a polypeptide expressed from a polynucleotide selected from the group as shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14, as compared to the normal tissue, is identified as cancerous.

Even another embodiment of the invention provides a method of diagnosing dysplasia and cancer. The amount of specific mRNA molecules is determined in a test sample of tissue suspected of being dysplastic or cancerous and in a human tissue which is normal. The mRNA molecules measured are complementary to the minus strand of a double-stranded polynucleotide sequence. The double-stranded polynucleotide sequence is selected from the group of polynucleotides as shown in SEQ ID NOs:2, 5, and 15. The determined amounts of mRNA molecules are compared. A

test sample of human tissue which contains more of the mRNA molecules than the normal tissue is identified as being dysplastic or cancerous.

Yet another embodiment of the invention provides a method of diagnosing dysplasia. The amounts of specific mRNA molecules in a test sample of human tissue suspected of being dysplastic and in a human tissue which is normal are determined. The mRNA molecules are complementary to the minus strand of a double-stranded polynucleotide sequence. The double-stranded polynucleotide sequence is selected from the group of polynucleotides as shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14. The determined amounts of mRNA molecules are then compared. A test sample of human tissue which contains more of the mRNA molecules than the normal tissue is identified as being dysplastic.

Still another embodiment of the invention provides a method of diagnosing cancer. The amounts of a first set of specific mRNA molecules in a test sample of tissue of a human suspected of being cancerous and in a human tissue which is normal are determined. The mRNA molecules are complementary to the minus strand of a double-stranded polynucleotide sequence. The double-stranded polynucleotide sequence is selected from the group of polynucleotide sequences as shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14. In addition, the amounts of a second set of specific mRNA molecules in a test sample of tissue of a human suspected of being cancerous and in a human tissue which is normal are determined. The mRNA molecules are complementary to the minus strand of a double-stranded polynucleotide sequence. The double-stranded polynucleotide sequence is selected from the group of polynucleotide sequences as shown in SEQ ID NOs:2, 5, and 15. The determined amounts of the first and second sets of mRNA molecules are compared. A test sample of human tissue which contains more of the second set of mRNA molecules than the normal tissue, and which contains substantially the same amount of the first set of mRNA molecules, as compared to the normal tissue, is identified as cancerous.

Yet another embodiment of the invention provides a therapeutic composition useful for decreasing the amount of translation of an mRNA molecule in a cell. The composition comprises an antisense polynucleotide complementary to the plus strand of a double-stranded polynucleotide. The double-stranded polynucleotide is

selected from the group consisting of polynucleotides comprising a nucleotide sequence as shown in SEQ ID NOs:1-11, and 13-15, wherein said antisense polynucleotide binds to an mRNA molecule. The composition also includes a pharmaceutically acceptable carrier.

5 A further embodiment of the invention provides a therapeutic composition useful for reducing the expression of a polypeptide. The composition comprises an antibody which specifically binds to a polypeptide expressed from a polynucleotide selected from the group consisting of polynucleotides comprising a nucleotide sequence as shown in SEQ ID NOs:1-11 and 13-15. The composition also
10 includes a pharmaceutically acceptable carrier.

Another embodiment of the invention provides a therapeutic composition useful for reducing the translation from an mRNA molecule. The composition comprises a ribozyme which binds to an mRNA molecule, wherein a portion of said ribozyme is complementary to the plus strand of a double-stranded
15 polynucleotide. The polynucleotide is selected from the group consisting of the polynucleotides comprising a sequence as shown in SEQ ID NOs:1-11, and 13-15. The composition also comprises a pharmaceutically acceptable carrier.

The present invention provides the art with useful polynucleotides which represent expressed sequences of genes. Expression of the genes is mis-regulated in
20 cancer. The invention also provides the art with diagnostic methods based on the over- and under-expression of the genes and the polypeptides encoded by the genes in cancer and dysplastic cells. Inhibitors of the over-expressed polynucleotides and polypeptides can be used to reduce the growth of cancer cells and dysplastic cells. The polynucleotides and polypeptides which are under-expressed in cancer and dysplasia
25 can be delivered therapeutically to reduce the abnormal characteristics of cancer cells and dysplastic cells.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Polynucleotides that are mis-regulated in cancer and dysplasia are disclosed. The mis-regulated polynucleotide sequences are shown in SEQ ID NOs:1-
30 15. The polynucleotides are mis-regulated as follows:

SEQ ID NO:12 is down-regulated in cancer;

SEQ ID NOs:2, 5, and 15 are up-regulated in cancer and dysplasia; and

SEQ ID NOs:1, 3-4, 6-11, and 13-14 are up-regulated in dysplasia only.

Polynucleotides that are differentially regulated in cancer or dysplasia or
5 both can be useful in the diagnosis and treatment of these diseases. Dysplasia is an atypical proliferation of epithelial or mesenchymal cells that may represent an early stage of cancer; however, dysplasia does not necessarily progress to cancer. Epithelial dysplasia results in the loss of normal orientation of one epithelial cell to another, accompanied by alterations in cellular and nuclear size and shape. Cancer is a
10 proliferation of malignant cells that are no longer under normal physiologic control.

The subgenomic polynucleotides of the invention contain less than a whole chromosome and are preferably intron-free. The subgenomic polynucleotides of the invention can be isolated and purified free from other nucleotide sequences by standard nucleic acid purification techniques, for example, using PCR, cloning, and/or
15 restriction enzymes and probes to isolate fragments comprising the encoding sequences. Subgenomic polynucleotides of the invention can include all or a contiguous portion of a gene coding region. In one embodiment, an isolated and purified subgenomic polynucleotide of the invention comprises at least 10, 11, 12, 15, 18, 20, 25, 30, 35, 40, 45, 50, 60, 70, 74, 80, 90, 100, 125, 150, 154, 175, 200, 250, 300, or 350 contiguous
20 nucleotides selected from the polynucleotide sequences as shown in SEQ ID NOs:1-15. In a preferred embodiment, the polynucleotide molecules comprise a contiguous sequence of at least twelve nucleotides selected from the group consisting of the polynucleotides shown in SEQ ID NOs:1-15.

An open reading frame is a region of DNA that consists exclusively of
25 triplets that represent amino acids. The open reading frame of the polynucleotide sequences of the invention can be determined by examining all three possible reading frames in both directions. If a reading frame contains termination codons it cannot be read into protein and is not considered an open reading frame. Usually, no more than one of the six possible frames is open in any single stretch of DNA. An extensive open
30 reading frame is unlikely to exist by chance because of the lack of selective pressure to prevent the accumulation of nonsense codons. Therefore, the identification of a lengthy

open reading frame is taken to be *prima facie* evidence that the sequence is translated into protein in that frame. Lewin, ed., 1990, *Genes IV*, Cell Press, Cambridge, Mass.

Subgenomic polynucleotides of the invention can be used, *inter alia*, to produce proteins or polypeptides, as probes for the detection of mRNA of the invention in samples or extracts of human cells, to generate additional copies of the polynucleotides, and to generate ribozymes or antisense oligonucleotides. The subgenomic polynucleotides can also be used as single stranded DNA probes or as triple-strand forming oligonucleotides. The probes can be used to determine the presence or absence of the polynucleotide sequences as shown in SEQ ID NOs:1-15 or variants thereof in a sample.

The sequence of a nucleic acid comprising at least 15 contiguous nucleotides of at least any one of SEQ ID NO:1-15, preferably the entire sequence of at least any one of SEQ ID NO:1-15, is not limited and can be any sequence of A, T, G, and/or C (for DNA) and A, U, G, and/or C (for RNA) or modified bases thereof, including inosine and pseudouridine. The choice of sequence will depend on the desired function and can be dictated by coding regions desired, the intron-like regions desired, and the regulatory regions desired.

Where the entire sequence of any one of SEQ ID NO:1-15 is within the nucleic acid, the nucleic acid obtained is referred to herein as a polynucleotide comprising the sequence of any one of SEQ ID NO:1-15.

Both secreted and membrane-bound polypeptides of the present invention are of interest. For example, levels of secreted polypeptides can be assayed conveniently in body fluids, such as blood and urine. Membrane-bound polypeptides are useful for constructing vaccine antigens or inducing an immune response. Such antigens would comprise all or part of the extracellular region of the membrane-bound polypeptides.

Because both secreted and membrane-bound polypeptides comprise a fragment of contiguous hydrophobic amino acids, hydrophobicity predicting algorithms can be used to identify such polypeptides.

A signal sequence is usually encoded by both secreted and membrane-bound polypeptide genes to direct a polypeptide to the surface of the cell. The signal

sequence usually comprises a stretch of hydrophobic residues. Such signal sequences can fold into helical structures.

Membrane-bound polypeptides typically comprise at least one transmembrane region that possesses a stretch of hydrophobic amino acids that can transverse the membrane. Some transmembrane regions also exhibit a helical structure.

Hydrophobic fragments within a polypeptide can be identified by using computer algorithms. Such algorithms include Hopp & Woods, *Proc. Natl. Acad. Sci. USA* 78:3824-3828 (1981); Kyte & Doolittle, *J. Mol. Biol.* 157:105-132 (1982); and RAOAR algorithm, Degli Esposti *et al.*, *Eur. J. Biochem.* 190:207-219 (1990).

Another method of identifying secreted and membrane-bound polypeptides is to translate the present polynucleotides, SEQ ID NO:1-15, in all six frames and determine if at least 8 contiguous hydrophobic amino acids are present. Those translated polypeptides with at least 8; more typically, 10; even more typically, 12 contiguous hydrophobic amino acids are considered to be either a putative secreted or membrane bound polypeptide. Hydrophobic amino acids include alanine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, threonine, tryptophan, tyrosine, and valine.

The polypeptides of the invention include those encoded by the disclosed polynucleotides. These polypeptides can also be encoded by nucleic acids that, by virtue of the degeneracy of the genetic code, are not identical in sequence to the disclosed polynucleotides. Thus, the invention includes within its scope nucleic acids comprising polynucleotides encoding a protein or polypeptide expressed by a polynucleotide having the sequence of any one of SEQ ID NO:1-15. Also within the scope of the invention are variants; variants of polypeptides include mutants, fragments, and fusions. Mutants can include amino acid substitutions, additions or deletions. The amino acid substitutions can be conservative amino acid substitutions or substitutions to eliminate non-essential amino acids, such as to alter a glycosylation site, a phosphorylation site or an acetylation site, or to minimize misfolding by substitution or deletion of one or more cysteine residues that are not necessary for function. Conservative amino acid substitutions are those that preserve the general charge, hydrophobicity/hydrophilicity, and/or steric bulk of the amino acid substituted. For

example, substitutions between the following groups are conservative: Gly/Ala, Val/Ile/Leu, Asp/Glu, Lys/Arg, Asn/Gln, Ser/Cys, Thr, and Phe/Trp/Tyr.

Cysteine-depleted muteins are variants within the scope of the invention.

These variants can be constructed according to methods disclosed in U.S. Patent No. 4,959,314, "Cysteine-Depleted Muteins of Biologically Active Proteins." The patent discloses how to substitute other amino acids for cysteines, and how to determine biological activity and effect of the substitution. Such methods are suitable for proteins according to this invention that have cysteine residues suitable for such substitutions, for example to eliminate disulfide bond formation.

10 The protein variants described herein are encoded by polynucleotides that are within the scope of the invention. The genetic code can be used to select the appropriate codons to construct the corresponding variants.

The invention encompasses polynucleotide sequences having at least 65% sequence identity to any one of SEQ ID NOs:1-15 as determined by the Smith-15 Waterman homology search algorithm as implemented in MSPRCH program (Oxford Molecular) using an affine gap search with the following search parameters: gap open penalty of 12, and gap extension penalty of 1.

Polynucleotide probes comprising at least 12 contiguous nucleotides selected from the nucleotide sequence of a polynucleotide of SEQ ID NO:1-15 are used 20 for a variety of purposes, including identification of human chromosomes and determining transcription levels.

The nucleotide probes are labeled, for example, with a radioactive, fluorescent, biotinylated, or chemiluminescent label, and detected by well known methods appropriate for the particular label selected. Protocols for hybridizing 25 nucleotide probes to preparations of metaphase chromosomes are also well known in the art. A nucleotide probe will hybridize specifically to nucleotide sequences in the chromosome preparations which are complementary to the nucleotide sequence of the probe. A probe that hybridizes specifically to a polynucleotide should provide a detection signal at least 5-, 10-, or 20-fold higher than the background hybridization 30 provided with other unrelated sequences.

Polynucleotides of the present invention are used to identify a chromosome on which the corresponding gene resides. Using fluorescence in situ hybridization (FISH) on normal metaphase spreads, comparative genomic hybridization allows total genome assessment of changes in relative copy number of DNA sequences.

- 5 See Schwartz and Samad, *Current Opinions in Biotechnology* (1994) 8:70-74; Kallioniemi *et al.*, *Seminars in Cancer Biology* (1993) 4:41-46; Valdes and Tagle, *Methods in Molecular Biology* (1997) 68:1; Boultonwood, ed., Human Press, Totowa, NJ.

Preparations of human metaphase chromosomes are prepared using standard cytogenetic techniques from human primary tissues or cell lines. Nucleotide probes comprising at least 12 contiguous nucleotides selected from the nucleotide sequence of SEQ ID NOs:1-15 are used to identify the corresponding chromosome. The nucleotide probes are labeled, for example, with a radioactive, fluorescent, biotinylated, or chemiluminescent label, and detected by well known methods appropriate for the particular label selected. Protocols for hybridizing nucleotide probes to preparations of metaphase chromosomes are also well known in the art. A nucleotide probe will hybridize specifically to nucleotide sequences in the chromosome preparations that are complementary to the nucleotide sequence of the probe. A probe that hybridizes specifically to a polynucleotide-related gene provides a detection signal at least 5-, 10-, or 20-fold higher than the background hybridization provided with non-polynucleotide coding sequences.

Polynucleotides are mapped to particular chromosomes using, for example, radiation hybrids or chromosome-specific hybrid panels. See Leach *et al.*, *Advances in Genetics*, (1995) 33:63-99; Walter *et al.*, *Nature Genetics* (1994) 7:22-28; Walter and Goodfellow, *Trends in Genetics* (1992) 9:352. Such mapping can be useful in identifying the function of the polynucleotide-related gene by its proximity to other genes with known function. Function can also be assigned to the related gene when particular syndromes or diseases map to the same chromosome.

A polynucleotide will be useful in forensics, genetic analysis, mapping, and diagnostic applications if the corresponding region of a gene is polymorphic in the human population. A particular polymorphic form of the polynucleotide may be used to either identify a sample as deriving from a suspect or rule out the possibility that the

sample derives from the suspect. Any means for detecting a polymorphism in a gene are used, including but not limited to electrophoresis of protein polymorphic variants, differential sensitivity to restriction enzyme cleavage, and hybridization to an allele-specific probe.

5 Any naturally occurring variants of the nucleotide sequences which encode variants thereof are within the scope of this invention. Allelic variants of subgenomic polynucleotides of the invention can occur and can be identified by hybridization of putative allelic variants with nucleotide sequences disclosed herein under stringent conditions. For example, by using the following wash conditions--2 x
10 SCC, 0.1% SDS, room temperature twice, 30 minutes each; then 2 x SCC, 0.1% SDS, 50 °C once, 30 minutes; then 2 x SCC, room temperature twice, 10 minutes each--allelic variants of the polynucleotides of the invention can be identified which contain at most about 25-30% base pair mismatches. More preferably, allelic variants contain 15-25% base pair mismatches, even more preferably 5-15%, or 2-5%, or 1-2% base pair
15 mismatches.

Amplification by the polymerase chain reaction (PCR) can be used to obtain the polynucleotides of the invention, using either genomic DNA or cDNA as a template. The polynucleotides of the invention may also be obtained using reverse transcriptase and mRNA molecules that are complementary to the minus strand of a
20 double-stranded sequence wherein said double-stranded sequence is selected from the group of polynucleotides comprising a sequence as shown in SEQ ID NOs:1-15. Using the polynucleotide sequences disclosed herein, subgenomic polynucleotide molecules of the invention can also be made using the techniques of synthetic chemistry.

Probes specific to the polynucleotides of the invention may be generated
25 using the polynucleotide sequences disclosed in SEQ ID NOs:1-15. The probes are preferably at least 12, 14, 16, 18, 20, 22, 24, or 25 nucleotides in length and can be less than 2, 1, 0.5, 0.1, or 0.05 kb in length. The probes can be synthesized chemically or can be generated from longer polynucleotides using restriction enzymes. The probes can be labeled, for example, with a radioactive, biotinylated, or fluorescent tag.

30 Subgenomic polynucleotides of the invention can be propagated in vectors and cell lines using techniques well known in the art. Expression systems in

bacteria include those described in Chang *et al.*, *Nature* (1978) 275: 615; Goeddel *et al.*, *Nature* (1979) 281: 544; Goeddel *et al.*, *Nucleic Acids Res.* (1980) 8: 4057; EP 36,776; U.S. 4,551,433; deBoer *et al.*, *Proc. Natl. Acad. Sci. USA* (1983) 80: 21-25; and Siebenlist *et al.*, *Cell* (1980) 20: 269.

- 5 Expression systems in yeast include those described in Hinnen *et al.*, *Proc. Natl. Acad. Sci. USA* (1978) 75: 1929; Ito *et al.*, *J. Bacteriol.* (1983) 153: 163; Kurtz *et al.*, *Mol. Cell. Biol.* (1986) 6: 142; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25: 141; Gleeson *et al.*, *J. Gen. Microbiol.* (1986) 132: 3459; Roggenkamp *et al.*, *Mol. Gen. Genet.* (1986) 202: 302; Das *et al.*, *J. Bacteriol.* (1984) 158: 1165; De Louvencourt *et al.*, *J. Bacteriol.* (1983) 154: 737; Van den Berg *et al.*, *Bio/Technology* (1990) 8: 135; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25: 141; Cregg *et al.*, *Mol. Cell. Biol.* (1985) 5: 3376; U.S. 4,837,148; US 4,929,555; Beach and Nurse, *Nature* (1981) 300: 706; Davidow *et al.*, *Curr. Genet.* (1985) 10: 380; Gaillardin *et al.*, *Curr. Genet.* (1985) 10: 49; Ballance *et al.*, *Biochem. Biophys. Res. Commun.* (1983) 112: 284-289; Tilburn *et al.*, *Gene* (1983) 26: 205-221; Yelton *et al.*, *Proc. Natl. Acad. Sci. USA* (1984) 81: 1470-1474; Kelly and Hynes, *EMBO J.* (1985) 4: 475479; EP 244,234; and WO 91/00357.
- 10
- 15

- Expression of the subgenomic polynucleotides of the invention in insects can be accomplished as described in U.S. 4,745,051, Friesen *et al.* (1986) "The Regulation of Baculovirus Gene Expression" in: THE MOLECULAR BIOLOGY OF BACULOVIRUSES (W. Doerfler, ed.); EP 127,839; EP 155,476; Vlak *et al.*, *J. Gen. Virol.* (1988) 69: 765-776; Miller *et al.*, *Ann. Rev. Microbiol.* (1988) 42: 177; Carbonell *et al.*, *Gene* (1988) 73: 409; Maeda *et al.*, *Nature* (1985) 315: 592-594; Lebacqz-Verheyden *et al.*, *Mol. Cell. Biol.* (1988) 8: 3129; Smith *et al.*, *Proc. Natl. Acad. Sci. USA* (1985) 82: 8404; Miyajima *et al.*, *Gene* (1987) 58: 273; and Martin *et al.*, *DNA* (1988) 7: 99. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts are described in Luckow *et al.*, *Bio/Technology* (1988) 6: 47-55; Miller *et al.*, in GENETIC ENGINEERING (Setlow, J.K. *et al.* eds.), Vol. 8 (Plenum Publishing, 1986), pp. 277-279; and Maeda *et al.*, *Nature*, (1985) 315: 592-594.
- 20
- 25

- 30 Mammalian expression of the subgenomic polynucleotides of the invention can be accomplished as described in Dijkema *et al.*, *EMBO J.* (1985) 4: 76;

Gorman *et al.*, *Proc. Natl. Acad. Sci. USA*, (1982) 79: 6777; Boshart *et al.*, *Cell* (1985) 41: 521; and U.S. 4,399,216. Other features of mammalian expression can be facilitated as described in Ham and Wallace, *Meth. Enz.* (1979) 58: 44; Barnes and Sato, *Anal. Biochem.* (1980) 102: 255; U.S. 4,767,704; US 4,657,866; US 4,927,762; US 4,560,655; WO 90/103430; WO 87/00195; and U.S. RE 30,985.

The subgenomic polynucleotides of the invention can be on linear or circular molecules. They can be on autonomously replicating molecules (vectors) or on molecules without replication sequences. They can be regulated by their own or by other regulatory sequences, as is known in the art. The subgenomic polynucleotides of the invention can be introduced into suitable host cells using a variety of techniques which are available in the art, such as transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated DNA transfer, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, gene gun, and calcium phosphate-mediated transfection.

The invention provides a method of detecting expression of a polynucleotide in, for example, a biological sample, which can be useful, *inter alia*, for diagnosing cancer or dysplasia. The basis for this method is the discovery that the polynucleotide sequence(s) as shown in:

SEQ ID NO:12 is down-regulated in cancer;

SEQ ID NOs:2, 5, and 15 are up-regulated in cancer and dysplasia; and
SEQ ID NOs:1, 3-4, 6-11, and 13-14 are up-regulated in dysplasia only.

In patients who have been diagnosed with pancreatic dysplasia or cancer, the detection of levels of the expression products of the polynucleotide sequences of the invention, either mRNA or protein, can be used to diagnose or prognose a disorder, to monitor treatment of the disorder, or to screen agents which affect the disorder.

The expression products of the polynucleotide sequences of the invention, either mRNA or proteins, can be detected in a body sample for diagnosis or prognosis. The body sample can be, for example, a solid tissue or a fluid sample. The patient from whom the body sample is obtained can be healthy or can already be identified as having a condition in which altered expression of a protein of the invention is implicated.

In one embodiment, the body sample is assayed for the level of a protein expressed from a polynucleotide sequence of the invention. The protein could be detected by, for example, antibodies to the proteins. The antibodies can be labeled, for example, with a radioactive, fluorescent, biotinylated, or enzymatic tag and detected directly, or can be detected using indirect immunochemical methods, using a labeled secondary antibody. The presence of the protein can be assayed, for example, in tissue sections by immunocytochemistry, or in lysates, using Western blotting, as is known in the art.

The level of the protein in a tissue sample suspected of being cancerous or dysplastic is compared with the level of the protein in a normal tissue. A higher level of the polypeptides expressed from polynucleotide sequences as shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14 in the suspect tissue, as compared to the normal tissue, indicates the presence of dysplastic cells in the suspect tissue. A higher level of the polypeptides expressed from polynucleotide sequences as shown in SEQ ID NOs:2, 5, and 15 in the suspect tissue, as compared to the normal tissue, indicates the presence of dysplastic cells or cancerous cells or both in the suspect tissue. A lower level of the polypeptide expressed from the polynucleotide sequence as shown in SEQ ID NO:12 in the suspect tissue, as compared to the normal tissue, indicates the presence of cancerous cells in the suspect tissue.

Additionally, a differentiation between cancer or dysplasia in a patient's diagnosis can be made. The expression of a polynucleotide sequence of the invention that is up-regulated in dysplastic cells only (*i.e.*, SEQ ID NOs: 1, 3-4, 6-11, and 13-14) and the expression of a polynucleotide that is up-regulated in both dysplastic cells and cancerous cells (*i.e.*, SEQ ID NOs:2, 5, and 15) can be used to screen a patient's tissues. If examination of a patient's tissues reveals that there is no up-regulation of a polynucleotide sequence that is up-regulated in dysplastic cells only (*i.e.*, SEQ ID NOs: 1, 3-4, 6-11, and 13-14), and that there is up-regulation of a polynucleotide sequence that is up-regulated in both cancerous cells and dysplastic cells (*i.e.*, SEQ ID NOs:2, 5, and 15), then the patient is diagnosed with cancer.

Alternatively, the presence of mRNA expressed from the polynucleotide sequences of the invention in two tissues can be compared. mRNA can be detected, for

example, by *in situ* hybridization in tissue sections, by reverse transcriptase-PCR, or in Northern blots containing poly A+ mRNA. One of skill in the art can readily determine differences in the size or amount of mRNA transcripts between two tissues, using Northern blots and nucleotide probes. For example, the level of mRNA of the invention
5 in a tissue sample suspected of being cancerous or dysplastic is compared with the expression of the mRNA in a normal tissue. Any methods known in the art for determining the amounts of specific mRNAs can be used.

A higher level of mRNA expressed from polynucleotide sequences as shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14 in the suspect tissue, as compared to the
10 normal tissue, indicates the presence dysplastic cells in the suspect tissue. A higher level of mRNA expressed from the polynucleotide sequences as shown in SEQ ID NOs:2, 5, and 15 in the suspect tissue, as compared to the normal tissue, indicates the presence dysplastic cells or cancerous cells or both in the suspect tissue. A lower level of the mRNA expressed from the polynucleotide sequence as shown in SEQ ID NO:12
15 in the suspect tissue, as compared to the normal tissue, indicates the presence of cancerous cells in the suspect tissue. Any combinations of these sequences can be used to determine a diagnosis.

Optionally, the level of a particular expression product of a polynucleotide sequence of the invention in a body sample can be quantitated.
20 Quantitation can be accomplished, for example, by comparing the level of expression product detected in the body sample with the amounts of product present in a standard curve. A comparison can be made visually or using a technique such as densitometry, with or without computerized assistance. Alternative methods can be used, for example ELISA, western blot, immunoprecipitation, radioimmunoassay, etc. Any method
25 known in the art for detecting and quantitating a particular protein can be used.

Reagents specific for the polynucleotides and polypeptides of the invention, such as antibodies and nucleotide probes, can be supplied in a kit for detecting the presence of an expression product in a biological sample. The kit can also contain buffers or labeling components, as well as instructions for using the reagents to
30 detect and quantify expression products in the biological sample.

Polynucleotide expression in a cell can be increased or decreased, as desired. Polynucleotide expression can be altered for therapeutic purposes, as described below, or can be used to identify and study the role of therapeutic agents in cancer and other diseases.

5 Decreasing the expression of genes containing sequences selected from the group consisting of the sequences as shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14 is useful, for example, as a therapeutic for altering the abnormal characteristics of dysplastic cells. Decreasing the expression of polynucleotide sequences selected from the group consisting of the sequences as shown in SEQ ID NOs:2, 5, and 15 is useful,
10 for example, as a therapeutic agent for decreasing the growth rate of dysplastic and cancer cells.

Expression of the polynucleotide sequences of the invention can be altered using an antisense oligonucleotide sequence. Therapeutic compositions for decreasing gene expression comprise an expression construct containing
15 polynucleotides encoding all or a portion of a polynucleotide sequence selected from the group consisting of SEQ ID NOs:1-11, and 13-15. Within the expression construct, the polynucleotide segment is orientated in the antisense direction and is located downstream from a promoter. Transcription of the polynucleotide segment initiates at the promoter.

20 Preferably, the antisense oligonucleotide sequence is at least ten nucleotides in length, but longer sequences of at least 11, 12, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 74, 80, 90, 100, 125, 150, 162, 175, 200, 250, 300, or 350 contiguous nucleic acids can also be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into cells whose division is to be decreased, as
25 described above. A more complete description of gene transfer vectors, especially retroviral vectors, is contained in U.S. Serial No. 08/869,309, which is incorporated herein by reference.

The antisense oligonucleotides can be composed of deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can
30 be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with phosphodiester or

non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamides, carboxymethyl esters, carbonates, and phosphate triesters. See Brown, 1994, *Meth. Mol. Biol.* 20:1-8; Sonveaux, 1994, *Meth. Mol. Biol.* 26:1-72; Uhlmann et al., 1990, *Chem. Rev.* 90:543-583.

Although precise complementarity is not required for successful duplex formation between an antisense molecule and the complementary coding sequence of a gene, antisense molecules with no more than one mismatch are preferred. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatch which will be tolerated between a particular antisense oligonucleotide and a particular coding sequence of the selected gene.

The antisense oligonucleotides of the invention can be modified without affecting their ability to hybridize to a polynucleotide coding sequence of the present invention. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside-phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases or sugars or both, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, can also be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art. Agrawal et al., 1992, *Trends Biotechnol.* 10:152-158; Uhlmann et al., 1990, *Chem. Rev.* 90:543-584; Uhlmann et al., 1987, *Tetrahedron. Lett.* 215:3539-3542.

Expression of the polynucleotides of the invention can also be decreased by delivering polyclonal, monoclonal, or single chain antibodies that specifically bind to polypeptides expressed from the polynucleotide sequences as shown in SEQ ID NOs:1-11 and 13-15. Antibodies specific to these proteins bind to the protein and prevent the protein from functioning in the cell. Blocking protein expression or function is useful for preventing, reducing the effects of, or curing cancer and dysplasia.

In one embodiment of the invention, expression of the polynucleotides selected from the group consisting of the polynucleotide sequences shown in SEQ ID

NOs: 1-11, and 13-15 are decreased using a ribozyme, an RNA molecule with catalytic activity. See, e.g., Cech, 1987, *Science* 236: 1532-1539; Cech, 1990, *Ann. Rev. Biochem.* 59:543-568; Cech, 1992, *Curr. Opin. Struct. Biol.* 2: 605-609; Couture and Stinchcomb, 1996, *Trends Genet.* 12: 510-515. Ribozymes can be used to inhibit gene
5 function by cleaving an RNA sequence, as is known in the art (e.g., Haseloff et al., U.S. 5,641,673).

The coding sequence of a polynucleotide of the invention can be used to generate a ribozyme which will specifically bind to RNA transcribed from said polynucleotide. Methods of designing and constructing ribozymes which can cleave
10 other RNA molecules in *trans* in a highly sequence specific manner have been developed and described in the art (see Haseloff, J. et al. (1988), *Nature* 334:585-591). For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically
15 hybridizes with the target (see, for example, Gerlach, W. L. et al., EP 321,201). Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme of the invention can be integrally related; thus, upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

20 Ribozymes of the invention can be introduced into cells as part of a DNA construct, as is known in the art. The DNA construct can also include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling the transcription of the ribozyme in the cells.

25 Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, gene gun, or calcium phosphate precipitation, can be used to introduce the ribozyme-containing DNA construct into cells whose division it is desired to decrease, as described above. Alternatively, if it is desired that the DNA construct be stably retained by the cells, the DNA construct can be supplied on a
30 plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art.

As taught in Haseloff et al., U.S. 5,641,673, the ribozymes of the invention can be engineered so that their expression will occur in response to factors which induce expression of a polynucleotide of the invention. The ribozyme can also be engineered to provide an additional level of regulation, so that destruction of RNA occurs only when both the ribozyme and the corresponding gene are induced in the cells.

Preferably, the mechanism used to decrease expression of the polynucleotide of the invention, whether antisense nucleotide sequence, antibody, or ribozyme decreases expression of the polynucleotide by 50%, 60%, 70%, or 80%. Most preferably, expression of the polynucleotide is decreased by 90%, 95%, 99%, or 100%. The effectiveness of the mechanism chosen to alter expression of the polynucleotide can be assessed using methods well known in the art, such as hybridization of nucleotide probes to mRNA of the polynucleotide, quantitative RT-PCR, or detection of a protein using specific antibodies of the invention.

Increased expression of a polynucleotide is useful to decrease the growth rate of cancer cells where the particular polynucleotide is down-regulated in cancer cells, such as the polynucleotide sequence as shown in SEQ ID NO:12. Therapeutic compositions for increasing polynucleotide expression comprise an expression construct containing all or a portion of the polynucleotide sequence as shown in SEQ ID NO:12. Within an expression construct, the polynucleotide segment is oriented in the sense direction and is located downstream from the promoter. Transcription of the polynucleotide segment initiates at the promoter. The expression construct can be introduced into cells along with a pharmaceutically acceptable carrier to decrease the growth rate of cancer cells or ameliorate other abnormal characteristics. Expression of the polynucleotide sequence can be monitored by detecting production of mRNA which hybridizes to the delivered polynucleotide or by detecting protein encoded by the delivered polynucleotide.

Proteins that are expressed from the polynucleotide sequences of the invention can be produced recombinantly in prokaryotic or eukaryotic host cells, such as bacteria, yeast, insect, or mammalian cells, using expression vectors known in the art. Enzymes can be used to generate less than full length polypeptides by enzymatic

proteolysis of full-length proteins of the invention. Alternatively, synthetic chemistry methods, such as solid-phase peptide synthesis, can be used to synthesize the proteins and polypeptides.

Species homologs of human subgenomic polynucleotides or the encoded polypeptides can be identified by making suitable probes or primers and screening cDNA expression libraries from other species, such as mice, monkeys, yeast, or bacteria. Mammalian homologs are preferred, however.

Proteins or polypeptides expressed from the polynucleotide sequences as shown in SEQ ID NO:1-15 can be isolated and purified from human cells that express the proteins. The proteins can be obtained substantially free from other human proteins by standard protein purification methods, such as size exclusion chromatography, ion exchange chromatography, ammonium sulfate fractionation, affinity chromatography, or preparative gel electrophoresis.

Proteins or polypeptides expressed from the polynucleotides of the invention can also be used in a fusion protein, for example, as an immunogen. The fusion protein comprises two protein segments. The first protein segment consists of at least six, eight, ten, twelve, fifteen, twenty or thirty contiguous amino acids of a polypeptide sequence expressed from a polynucleotide sequence as shown in SEQ ID NOs:1-15. The first protein segment is fused to a second protein segment by means of a peptide bond. The second protein segment can be a full-length protein or a fragment of a protein. Techniques for making fusion proteins, either recombinantly or by covalently linking two protein segments, are well known in the art.

The second protein or protein fragment of a fusion protein can be derived from another type of protein or a similar protein. The second protein or protein fragment can be labeled with a detectable marker, such as a radioactive or fluorescent tag, or can be an enzyme that will generate a detectable product. Enzymes suitable for this purpose, such as β -galactosidase, are well-known in the art. A fusion protein can be used, for example, to target the proteins of the invention or polypeptides to a particular location in a cell or tissue, in various assays, such as the yeast two-hybrid technique, or as an immunogen.

The proteins or polypeptides expressed from the polynucleotides of the invention can be used for generating antibodies. The antibodies can be used, *inter alia*, to detect and quantitate expression of the cognate protein. Proteins or polypeptides expressed from the polynucleotides of the invention comprising at least six, eight, ten, 5 twelve, fifteen, twenty or thirty consecutive amino acids can be used as immunogens. The proteins or polypeptides can be used to obtain a preparation of antibodies which specifically bind to a protein or polypeptide of the invention. The antibodies can be polyclonal or monoclonal. Techniques for raising both polyclonal and monoclonal antibodies are well known in the art.

10 Single chain antibodies can also be constructed. Single chain antibodies which specifically bind to a protein or polypeptide expressed from the polynucleotides of the invention can be isolated, for example, from single-chain immunoglobulin display libraries, as are known in the art. The library is "panned" against a protein or polypeptide, and a number of single chain antibodies which bind different epitopes of 15 the polypeptide with high-affinity can be isolated. Hayashi *et al.*, 1995, *Gene* 160:129-30. Such libraries are known and available to those in the art. The antibodies can also be constructed using the polymerase chain reaction (PCR), using hybridoma cDNA as a template. Thirion *et al.*, 1996, *Eur. J. Cancer Prev.* 5:507-11.

The single chain antibody can be mono- or bi-specific, and can be 20 bivalent or tetravalent. Construction of tetravalent bispecific single chain antibodies is taught in Coloma and Morrison, 1997, *Nat. Biotechnol.* 15:159-63. Construction of bivalent bispecific single chain antibodies is taught in Mallender and Voss, 1994, *J. Biol. Chem.* 269:199-206.

A nucleotide sequence encoding the single chain antibody can then be 25 constructed using manual or automated nucleotide synthesis, cloned into DNA expression vectors using standard recombinant DNA methodologies, and introduced into cells which express the selected gene, as described below. Alternatively, the antibodies can be produced directly using filamentous phage technology. Verhaar *et al.*, 1995, *Int. J. Cancer* 61:497-501; Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165:81-30 91.

The antibodies bind specifically to the epitopes of the proteins or polypeptides expressed from the polynucleotides of the invention. In a preferred embodiment, the epitopes are not present on other human proteins. Typically a minimum number of contiguous amino acids to encode an epitope is 6, 8, or 10.

5 However, more can be used, for example, at least 15, 25, or 50, especially to form epitopes which involve non-contiguous residues or particular conformations.

Antibodies that bind specifically to the proteins or polypeptides include those that bind to full-length proteins or polypeptides. Specific binding antibodies do not detect other proteins on Western blots of human cells, or provide a signal at least
10 ten-fold lower than the signal provided by the target protein of the invention. Antibodies which have such specificity can be obtained by routine screening. In a preferred embodiment of the invention, the antibodies immunoprecipitate the proteins or polypeptides expressed from the polynucleotides of the invention from cell extracts or solution. Additionally, the antibodies can react with proteins or polypeptides
15 expressed from the polynucleotides of the invention in tissue sections or on Western blots of polyacrylamide gels. Preferably the antibodies do not exhibit nonspecific cross-reactivity with other human proteins on Western blots or in immunocytochemical assays.

Techniques for purifying antibodies to the proteins or polypeptides
20 expressed from the polynucleotides of the invention are available in the art. In a preferred embodiment, the antibodies are passed over a column to which a particular protein or polypeptide expressed from the polynucleotides of the invention is bound. The bound antibodies are then eluted, for example, with a buffer having a high salt concentration.

25 Therapeutic compositions of the invention also comprise a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers are well known to those in the art. Such carriers include, but are not limited to, large, slowly metabolized macromolecule, such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and inactive virus
30 particles. Pharmaceutically acceptable salts can also be used in the composition, for example, mineral salts such as hydrochlorides, hydrobromides, phosphates, or sulfates,

as well as the salts of organic acids such as acetates, propionates, malonates, or benzoates.

Therapeutic compositions can also contain liquids, such as water, saline, glycerol, and ethanol, as well as substances such as wetting agents, emulsifying agents, or pH buffering agents. Liposomes, such as those described in U.S. 5,422,120, WO 95/13796, WO 91/14445, or EP 524,968 B1, can also be used as a carrier for the therapeutic composition.

Typically, a therapeutic composition is prepared as an injectable, either as a liquid solution or suspension; however, solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection can also be prepared. A composition can also be formulated into an enteric coated tablet or gel capsule according to known methods in the art, such as those described in U.S. 4,853,230, EP 225,189, AU 9,224,296, and AU 9,230,801.

Administration of the therapeutic agents of the invention can include local or systemic administration, including injection, oral administration, particle gun, or catheterized administration, and topical administration. Various methods can be used to administer a therapeutic composition directly to a specific site in the body.

For treatment of tumors, for example, a small tumor or metastatic lesion can be located and a therapeutic composition injected several times in several different locations within the body of the tumor. Alternatively, arteries which serve a tumor can be identified, and a therapeutic composition injected into such an artery, in order to deliver the composition directly into the tumor.

A tumor which has a necrotic center can be aspirated and the composition injected directly into the now empty center of the tumor. A therapeutic composition can be directly administered to the surface of a tumor, for example, by topical application of the composition. X-ray imaging can be used to assist in certain of the above delivery methods. Combination therapeutic agents, including a protein or polypeptide or a subgenomic polynucleotide and other therapeutic agents, can be administered simultaneously or sequentially.

Receptor-mediated targeted delivery can be used to deliver therapeutic compositions containing subgenomic polynucleotides, proteins, or reagents such as

antibodies, ribozymes, or antisense oligonucleotides of the invention to specific tissues. Receptor-mediated delivery techniques are described in, for example, Findeis et al. (1993), *Trends in Biotechnol.* 11, 202-05; Chiou et al. (1994), *GENE THERAPEUTICS: METHODS AND APPLICATIONS OF DIRECT GENE TRANSFER* (J.A. Wolff, ed.); Wu & Wu
5 (1988), *J. Biol. Chem.* 263, 621-24; Wu et al. (1994), *J. Biol. Chem.* 269, 542-46; Zenke et al. (1990), *Proc. Natl. Acad. Sci. U.S.A.* 87, 3655-59; Wu et al. (1991), *J. Biol. Chem.* 266, 338-42.

Alternatively, therapeutic compositions can be introduced into human cells *ex vivo*, and the cells then replaced into the human. Cells can be removed from a
10 variety of locations including, for example, from a selected tumor or from an affected organ. In addition, a therapeutic composition can be inserted into non-affected, for example, dermal fibroblasts or peripheral blood leukocytes. If desired, particular fractions of cells such as a T cell subset or stem cells can also be specifically removed from the blood (*see*, for example, PCT WO 91/16116). The removed cells can then be
15 contacted with a therapeutic composition utilizing any of the above-described techniques, followed by the return of the cells to the human, preferably to or within the vicinity of a tumor or other site to be treated. The methods described above can additionally comprise the steps of depleting fibroblasts or other non-contaminating tumor cells subsequent to removing tumor cells from a human, and/or the step of
20 inactivating the cells, for example, by irradiation.

Both the dose of a composition and the means of administration can be determined based on the specific qualities of the therapeutic composition, the condition, age, and weight of the patient, the progression of the disease, and other relevant factors. Preferably, a therapeutic composition of the invention increases or decreases expression
25 of a polynucleotide by 50%, 60%, 70%, or 80%. Most preferably, expression of the polynucleotide is increased or decreased by 90%, 95%, 99%, or 100%. The effectiveness of the mechanism chosen to alter expression of the polynucleotide can be assessed using methods well known in the art, such as hybridization of nucleotide probes to mRNA of the polynucleotide, quantitative RT-PCR, or detection of a protein
30 or polypeptide using specific antibodies.

If the composition contains protein, polypeptide, or antibody, effective dosages of the composition are in the range of about 5 μ g to about 50 μ g/kg of patient body weight, about 50 μ g to about 5 mg/kg, about 100 μ g to about 500 μ g/kg of patient body weight, and about 200 to about 250 μ g/kg.

5 Therapeutic compositions containing subgenomic polynucleotides can be administered in a range of about 100 ng to about 200 mg of DNA for local administration in a gene therapy protocol. Concentration ranges of about 500 ng to about 50 mg, about 1 μ g to about 2 mg, about 5 μ g to about 500 μ g, and about 20 μ g to about 100 μ g of DNA can also be used during a gene therapy protocol. Factors such as
10 method of action and efficacy of transformation and expression are considerations that will effect the dosage required for ultimate efficacy of the subgenomic polynucleotides. Where greater expression is desired over a larger area of tissue, larger amounts of subgenomic polynucleotides or the same amounts re-administered in a successive protocol of administrations, or several administrations to different adjacent or close
15 tissue portions of, for example, a tumor site, may be required to effect a positive therapeutic outcome. In all cases, routine experimentation in clinical trials will determine specific ranges for optimal therapeutic effect.

 The therapeutic compositions are useful in treating pancreatic cancer and pancreatic dysplasia, as well as other types of cancers such as: bone cancer; brain
20 tumors; breast cancer; endocrine system cancers, such as cancers of the thyroid, pituitary, and adrenal glands and the pancreatic islets; gastrointestinal cancers, such as cancer of the anus, colon, esophagus, gallbladder, stomach, liver, and rectum; genitourinary cancers such as cancer of the penis, prostate and testes; gynecological cancers, such as cancer of the ovaries, cervix, endometrium, uterus, fallopian tubes,
25 vagina, and vulva; head and neck cancers, such as hypopharyngeal, laryngeal, oropharyngeal cancers, lip, mouth and oral cancers, cancer of the salivary gland, cancer of the aerodigestive tract and sinus cancer; leukemia; lymphomas including Hodgkin's and non-Hodgkin's lymphoma; metastatic cancer; myelomas; sarcomas; skin cancer; urinary tract cancers including bladder, kidney and urethral cancers; and pediatric
30 cancers, such as pediatric brain tumors, leukemia, lymphomas, sarcomas, liver cancer and neuroblastoma and retinoblastoma.

The following example provides data and experimental procedures. However, the invention is not limited to the example. The invention is defined in the specification as a whole which includes the claims.

EXAMPLE 1

5 A family was identified that had several members who had been diagnosed with pancreatic cancer. The pathological features of disease in the family included progression from normal to metaplasia to dysplasia to cancer. Tissues were obtained from a member of the family diagnosed with pancreatic cancer, from a member of the family diagnosed with dysplasia of pancreatic cells, from a person
10 unrelated to the family diagnosed with pancreatitis, and from a person unrelated to the family with a normal pancreas.

Ductal cells from the tissues of each of these subjects were cultured and mRNA was isolated from the cultures. The mRNA was subjected to reverse transcriptase polymerase chain reaction using 200 primer pairs (10 anchored and 20
15 arbitrary primers). The resulting cDNA was subjected to a differential display in which the cDNA from each of the 4 samples were compared on a gel. Bands of cDNA that appeared to be up-or down-regulated in the dysplastic or pancreatic cancer samples, as compared to the normal and pancreatitis samples, were cut from the gel, amplified, cloned, and sequenced.

20 The following polynucleotides sequences, as shown in SEQ ID NOs: 1-15, were identified as being mis-regulated in pancreatic cancer or dysplasia or both:

TABLE 1

Up-Regulated and Down-Regulated Polynucleotides in Pancreatic Cancer and Dysplasia

<u>SEQ ID NO.</u>	<u>Regulation Status</u>
SEQ ID NO:1	Up in dysplasia only

<u>SEQ ID NO.</u>	<u>Regulation Status</u>
SEQ ID NO:2	Up in dysplasia and cancer
SEQ ID NO:3	Up in dysplasia only
SEQ ID NO:4	Up in dysplasia only
SEQ ID NO:5	Up in dysplasia and cancer
SEQ ID NO:6	Up in dysplasia only
SEQ ID NO:7	Up in dysplasia only
SEQ ID NO:8	Up in dysplasia only
SEQ ID NO:9	Up in dysplasia only
SEQ ID NO:10	Up in dysplasia only
SEQ ID NO:11	Up in dysplasia only
SEQ ID NO:12	Down in cancer only
SEQ ID NO:13	Up in dysplasia only
SEQ ID NO:14	Up in dysplasia only
SEQ ID NO:15	Up in dysplasia and cancer

All patents, published patent applications and publications cited herein are incorporated by reference as if set forth fully herein.

Although certain preferred embodiments have been described herein, it is not intended that such embodiments be construed as limitations on the scope of the

5 invention except as set forth in the following claims.

CLAIMS

We claim:

1. An isolated subgenomic polynucleotide comprising at least twelve contiguous nucleotides selected from the group consisting of the polynucleotide sequences as shown in SEQ ID NOs:1-15.
2. An isolated subgenomic polynucleotide comprising at least twelve contiguous nucleotides selected from the group consisting of the polynucleotide sequences of SEQ ID NOs:2, 5 and 15.
3. An isolated polypeptide, wherein said polypeptide comprises at least 6 contiguous amino acids encoded by a polynucleotide selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:1-15.
4. An isolated polypeptide, wherein said polypeptide comprises at least 6 contiguous amino acids encoded by a polynucleotide selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:2, 5 and 15.
5. An antibody preparation which specifically binds to a polypeptide of claim 3.
6. An antibody preparation which specifically binds to a polypeptide of claim 4.
7. An isolated nucleotide probe consisting of a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:1-15.
8. The polynucleotide of claim 1 wherein said polynucleotide sequence in SEQ ID NO:12.

9. A method of diagnosing cancer comprising the steps of:
determining the amount of a polypeptide expressed from the polynucleotide of claim 8 in a tissue sample of a human suspected of being cancerous and in a human tissue which is normal; and
comparing the determined amounts; wherein a sample human tissue which contains less of the polypeptide than the normal tissue is identified as cancerous.

10. A method of diagnosing cancer comprising the steps of:
determining the amount of mRNA molecules in a sample tissue of a human suspected of being cancerous and in a human tissue which is normal, wherein said mRNA molecules are complementary to the polynucleotide sequence of claim 8 or its complement; and
comparing the determined amounts of mRNA molecules; wherein a sample human tissue which contains less of said mRNA molecules than the normal tissue is identified as cancerous.

11. A therapeutic composition useful for reducing the growth rate of cancer cells comprising:
a polynucleotide comprising all or a portion of a nucleotide sequence of claim 8 which is operably linked to a promoter sequence, wherein said portion comprises at least 18 contiguous nucleotides; and
a pharmaceutically acceptable carrier.

12. A therapeutic composition useful for reducing the growth rate of cancer cells comprising:
a polypeptide comprising all or a portion of an amino acid sequence expressed from a polynucleotide of claim 8; and
a pharmaceutically acceptable carrier.

13. A method of diagnosing dysplasia and cancer comprising the steps of:
determining the amount of a polypeptide expressed from the polynucleotide of claim 2 in a sample tissue of a human suspected of being dysplastic or cancerous and in a human tissue which is normal; and
comparing the determined amounts; wherein a proband human tissue which contains more of the polypeptide than the normal tissue is identified as being dysplastic or cancerous.

14. A method of diagnosing dysplasia comprising the steps of:
determining the amount of a polypeptide expressed from a polynucleotide having a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14 in a sample human tissue suspected of being dysplastic and in a human tissue which is normal; and
comparing the determined amounts; wherein a sample human tissue which contains more of the polypeptide than the normal tissue is identified as being dysplastic.

15. A method of diagnosing cancer comprising the steps of:
determining the amount of a polypeptide expressed from a polynucleotide of claim 2, and the amount of a polypeptide expressed from a polynucleotide having a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14 in a sample human tissue suspected of containing cancer, and in a human tissue which is normal; and
comparing the determined amounts; wherein a sample human tissue which contains more of the polypeptide expressed from a polynucleotide of claim 2, as compared to the normal tissue and which contains substantially the same amount of a polypeptide expressed from a polynucleotide having a sequence selected from the group consisting of the polynucleotide sequences shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14, as compared to the normal tissue, is identified as cancerous.

16. A method of diagnosing dysplasia and cancer comprising the steps of:
determining the amount of mRNA molecules in a sample tissue of a human suspected of being dysplastic or cancerous and in a human tissue which is normal, wherein said mRNA molecules are complementary to the polynucleotide of claim 2 or its complement; and

comparing the determined amounts of mRNA molecules; wherein a sample human tissue which contains more of the mRNA molecules than the normal tissue is identified as being dysplastic or cancerous.

17. A method of diagnosing dysplasia comprising the steps of:
determining the amount of mRNA molecules in a sample human tissue suspected of being dysplastic and in a human tissue which is normal, wherein said mRNA molecules are complementary to the minus strand of a double-stranded sequence, wherein said double-stranded polynucleotide sequence is selected from the group of polynucleotides as shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14; and
comparing the determined amounts of mRNA molecules; wherein a sample human tissue which contains more of the mRNA molecules than the normal tissue is identified as being dysplastic.

18. A method of diagnosing cancer comprising the steps of:
determining the amount of a first set of mRNA molecules in a sample human tissue suspected of being cancerous and in a human tissue which is normal, wherein said mRNA molecules are complementary to the minus strand of a double-stranded polynucleotide sequence, wherein said double-stranded polynucleotide sequence is selected from the group of polynucleotides as shown in SEQ ID NOs:1, 3-4, 6-11, and 13-14; and
determining the amount of a second set of mRNA molecules in a sample human tissue suspected of being cancerous and in a human tissue which is normal, wherein said mRNA molecules are complementary to the minus strand of a double-stranded sequence, wherein said double-stranded polynucleotide sequence is selected from the group of polynucleotides as shown in SEQ ID NOs:2, 5, and 15; and

comparing the determined amounts; wherein a proband human tissue which contains more of the second set of mRNA molecules than the normal tissue, and which contains substantially same amount of the first set of mRNA molecules, as compared to the normal tissue, is identified as cancerous.

19. A therapeutic composition useful for decreasing the amount of translation of an mRNA molecule in a cell comprising:

an antisense polynucleotide complementary to the plus strand of a double-stranded polynucleotide, wherein said double-stranded polynucleotide comprises the polynucleotide of claim 2, wherein said antisense polynucleotide binds to an mRNA molecule; and

a pharmaceutically acceptable carrier.

20. A therapeutic composition useful for reducing the expression of a polypeptide comprising:

an antibody which specifically binds to a polypeptide expressed from the polynucleotide of claim 2; and

a pharmaceutically acceptable carrier.

21. A therapeutic composition useful for reducing the translation from an mRNA molecule comprising:

a ribozyme which binds to an mRNA molecule, wherein a portion of said ribozyme is complementary to the plus strand of a double-stranded polynucleotide;

wherein said polynucleotide comprises the polynucleotide of claim 2; and

a pharmaceutically acceptable carrier.

SEQUENCE LISTING

(1) GENERAL INFORMATION

(I) APPLICANT: KENNEDY, GIULIA

(ii) TITLE OF THE INVENTION: PANCREATIC CANCER GENES

(iii) NUMBER OF SEQUENCES: 15

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Seed and Berry LLP

(B) STREET: 701 Fifth Avenue

(C) CITY: Seattle

(D) STATE: WA

(E) COUNTRY: USA

(F) ZIP: 98104

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

(B) COMPUTER: IBM Compatible

(C) OPERATING SYSTEM: DOS

(D) SOFTWARE: FastSEQ for Windows Version 2.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 60/118570

(B) FILING DATE: June 3, 1999

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Potter, Jane E.R.

(B) REGISTRATION NUMBER: 33,332

(C) REFERENCE/DOCKET NUMBER: 200130.454

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 206-622-4900

(B) TELEFAX: 206-681-6031

(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(I) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 492 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGAACTCGG	TTTAAGACAG	GGCTTCTTCA	CCATTGCGAG	AGCGTTCACC	GGGACGAGTG	60
GCAAGAGTCT	TGGCTTGGAT	AGCATGAAGA	GCCCCAGTAC	AAGGAAGAAT	ACTGGAAATG	120
CTCAATTCGT	GGAGCGCGTT	TAAACGACGA	TTTATTTGGT	TTTCAATGAC	CGAGGACTTA	180
TGACAGGATG	ATTACATTTG	ACCTTGGGAC	ATGAACGCTT	GGACTGCTGA	CTTGTGTGTA	240
AAGCTGTTTT	GTTTGTGTGT	GTCTTGCTTG	ACAGTGGTTC	TCGATCATGG	TGATACCTGA	300
TGCTTTGGAC	ATGTCCACTT	ACTCCTCTAT	TATTCGTTGG	ATCATTGTTT	ATTCTGATAG	360
ATAGTGAATT	ATGTTCGGAT	GTGATCACA	GGATTGTGAT	TGTTAGTCCA	CTGTATCTCT	420
GATCGAATAG	GTCTATATAT	TATTATTTAG	ATAGAAAAAG	TAGCAATCCA	CTTAGGAGAT	480
TTATTGATCT	GC					492

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 545 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TCAGGTTTGA	GGCTGGAAAA	AGAATCATCC	CTTCCTTTCG	AGTTGAGATT	GTTTCTCATT	60
TTATAAGTAG	CTTTTATTTT	ATTTGAAATT	TGAATTTCTC	TTAAAATGGT	AGAGTATACC	120
AACCTTTACAG	AAAGGGGAAA	AAAGTCACCT	ACTGACTGAA	CACAGCTTTT	ACCAATTTGA	180
GCGTCTCCTT	GCAGTCTTTT	GAAATACGTA	TATGGGTAC	ACCATTGTAA	ACATGTGTTC	240
AGAGCTTGCA	ATTCATAAAT	ATGTTTATGT	CCGTTATCTA	ATGTGAGCTC	AAAACACAAT	300
AAGAGGGTCA	GGGTTGTGAA	GAAGGCAGGA	CAGGAATTAT	TTAACCATT	TTTCAAATGA	360
GAAAACCTGTG	GCCCAGATAC	AGAATGTCAC	TTGCTAAAT	CACATACATT	GAAACCAGTT	420
CTCTCCAGCA	TGTCACAGTG	CTTCTGTGTT	AGAGCCCAAG	TTACAAACCA	AAGTGTACAA	480
GGGCACAGAT	TATTAGCAAT	TTACATTTAA	AAATTTTAT	ATTTCCCTAAC	TGATACATAT	540
TAATT						545

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 978 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GTAGGTGTTG	TATTTCTACT	TTACAGGTAG	GAAAATGGAG	GCTAAGAAAA	GTTAATTTGT	60
CCGAGGGCCC	TCTGATGATA	GTGAAACTGG	GATGGAACTT	CTGCCTGCTT	GCTTCTGAGG	120
TCTGGGCTCC	TAACACTGTC	TCTACTGCCT	CGAGCCAAGA	GATTTACGCC	CTATTAAGCA	180
ATTTGTTGTG	CGATAAATTG	GAAGACACAG	CAGATAAGCA	AACAACTCAA	GCAACCAGGT	240
CGGTTCTCTG	AGTTTCTGAA	TTGTTGGGAC	CAAGGGGCCG	TGCAGAGGTA	ACCACAGCTG	300
GCGTAGTGTG	GTTGAGGTAG	CCCTATTAGC	CTTTTAGTTG	CTGTTACTAA	TTTATTTCTC	360
AGTGCTCAAT	GAACCAATTG	GCCATCAATC	AGCTTTGTGT	ATAGGTCATG	CTCCCATGGC	420
TCTGACCCAG	GTTGCTGCTC	AGAGTTGGCA	TCGTGGCTAA	AATATTACTA	GAGGTCAAAG	480
ATATGTGTGT	GTTTGTGGTT	GATTTAGTCG	AGTGATCTAG	AGGAATCTGA	ACCTTAGAGA	540
CTGAAGAAGA	ACCAGCATTT	CTGGGCAATA	ATACTTGAGT	TAAGGAGAGT	GTAGCAAAAC	600

TCTAGGTTAG	CATTGGCAGT	CCCTAGGATT	CAGACTGTAG	GCCTAAATGA	CCCTCAGTCC	660
AGAGCTGTAC	CTAATGAGGA	CAATACATTT	TAATGTGAGT	CCATTCTTAA	CAGCAAAATT	720
TCCTCTTTGC	TTGTCAACCAG	GGAAAAATGG	GTTTGCATAG	AAAAGGTGGA	GATTGAGGGG	780
GAAGCAGAAT	GGACAAGGAG	TAAAGAGGGA	ATCCAACACTAC	TTAGATTGTA	GCTTTCGTTC	840
TTCTTTGGTA	GTTGTAGAGG	TGAGCTTACC	AAAGCATAGA	TGACAGGCAA	TGTGGTATAC	900
AAGTTACTAC	ACTCCAAAAG	TCTGGGGTTC	TTACTTATTT	TGTGCATGAC	ATCCAAAGTA	960
GCCTAATAAA	ATCTTTTC					978

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 978 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GTAGATGTTG	TATTTCTACT	TTACAGGTAG	GAAAATGGAG	GCTAAGAAAA	GTTAATTTGT	60
CCGAGGGCCC	TCTGATGATA	GTGAACTGG	GATGGAACCT	CTGCCTGCTT	GCTTCTGAGG	120
TCTGGGCTCC	TAATACTGC	TCTACTGCCT	CGAGCCAAGA	GATTTACGCC	CTATTAAGCA	180
ATTTGTTGTG	CGATAAATTG	GAAGACACAG	CAGATAAGCA	AACAACCTCA	GCAACCAGGT	240
CAGTTCCTGG	AGTTTCTGAA	TTGTTGGGAC	CAAGGGGCCG	TGCAGAGGTA	ACCACAGCTG	300
GCGTAGTGTG	GTTGAGGTAG	CCCTATTAGC	CTTTTAGTTG	CTGTTACTAA	TTTATTTCTC	360
AGTGGTCAAT	GAACCAATTG	GCCATCAATC	AGCTTTGTGT	ATAGGTCATG	TTCCCATGGC	420
TCTGACCCAG	GTTGCTGCTC	AGAGTTGGCA	TCGTGGCTAA	AATATTACTA	GAGGTCAAAG	480
ATATGTGTGT	GTTTGTGGTT	GATTTAGTCG	AGTGATCTAG	AGGAATCTGA	ACCTTAGAGA	540
CTGAAGAAGA	ACCAGCATT	CTGGGCAATA	ATACTTGAGT	TAAGGAGAGT	GTAGCAAAAC	600
TCTAGGTTAG	CATTGGCAGT	CCCTAGGATT	CAGACTGTAG	GCCTAAATGA	CCCTCAGTCC	660
AGAGCTGTAC	CTAATGAGGA	CAATACATTT	TAATGTGAGT	CCATTCTTAA	CAGCAAAATT	720
TCCTCTTTGC	TTGTCAACCAG	GGAAAAATGG	GTTTGCATAG	AAAAGGTGGA	GATTGAGGGG	780
GAAGCAGAAT	GGACAAGGAG	TAAAGAGGGA	ATCCAACACTAC	TTAGATTGTA	GCTTTCGTTC	840
TTCTTTGGTA	GTTGTAGAGG	TGAGCTTACC	AAAGCATAGA	TGACAGGCAA	TGTGGTATAC	900
AAGTTACTAC	ACTCCAAAAG	TCTGGGGTTC	TTACTTATTT	TGTGCATGAC	ATCCAAAGTA	960
GCCTAATAAA	ATCTTTTC					978

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 539 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AATAGACATT	ATACTTTCTA	TGTGTGAAA	AGAGTTTTTC	AAAGATATGA	AACTGTAAAA	60
TATTTGTTAG	TTCCAGCCTA	TATATTTGCT	GGTTGGAGTA	TAGCTGACTC	ATTGAAATCA	120
AAGTCAATTT	TTTGAATTT	AATGTTTTTC	ATATGCTTGT	TCAGTGTTAT	AGTTCCTCAG	180
AAACTGCTGG	AATTTCTGTA	CTTCATTTTA	CCTTATGTCA	TTTATAGGCT	TAACATACCT	240
CTGCCTCCCA	CATCCAGACT	CATTTGTGAA	CTGAGCTGCT	ATGCAGTTGT	TAATTTTATA	300
ACTTTTTTCA	TCTTCTGAA	CAAGACTTTT	CAGTGGCCAA	ATAGTCAGGA	CATTCAAAGG	360
TTTATGTGGT	AATATCAGTG	ATATTTGCAA	CTGTGAAAAT	GGACTTAATA	ATTAGACCAT	420

TTCTACAAAG AACAACTGAA TAGGTGGAAG ACATGGAATT TCTTTTAGGT GCAGTGGTGG 480
TCTTCAAATT ACATTAGTTT TTTTATATA TATTTTAAAC ATATGTAAGA AATTAAGTG 539

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 491 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ATGAACTCGG TTTAAGACAG GGCTTCTTCA CCATTGCGAG AACGTTCCACC GGGACGAGTG 60
GCAAGAGTCT TGGCTTGGAT AGCATGAAGA GCCCCAGTAC AAGGAAGAAT ACTGGAAATG 120
CTCAATTCGT GGAGCGCGTT TAAACGACGA TTTATTTGGT TTTCAATGAC CGAGGACTTA 180
TGACAGGATG ATTACATTTG ACCTTGGGAC ATGAACGCTT GGACTGCTGA CTTGTGTGTA 240
AAGCTGTTTG TTTGTTTGTG TCTTGCTTGA CAGTGGTTCT CGATCATGAT GATACCTGAT 300
GCTTTGGACA TGTCCTACTTA CTCCTCTATT ATTCGTTGGA TCATTGTTTA TTCTGATAGA 360
TAGTGACTTA TGTTCCGATG TCGATCACAG GATTGTGATT GTTAGTCCAC TGTATCTCTG 420
ATCGAATAGG TCTATATATT ATTATTTAGA TAGAAAAAGT AGCAATCCAC TTAGGAGATT 480
TATTGATCTG C 491

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 491 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATGAACTCGG TTTAAGACAG GGCTTCTTCA CCATTGCGAG AACGTTCCACC GGGACGAGTG 60
GCAAGAGTCT TGGCTTGGAT AGCATGAAGA GCCCCAGTAC AAGGAAGAAT ACTGGAAATG 120
CTCAATTCGT GGAGCGCGTT TAAACGACGA TTTATTTGGT TTTCAATGAC CGAGACTTAT 180
GACAGGATGA TTACATTTGA CCTTGGGACA TGAACGCTTG GACTGCTGAC TTGTGTGTAA 240
AGCTGTTTTG TTTGTTTGTG TCTTGCTTGA CAGTGGTTCT CGATCATGAT GATACCTGAT 300
GCTTTGGACA TGTCCTACTTA CTCCTCTATT ATTCGTTGGA TCATTGTTTA TTCTGATAGA 360
TAGTGACTTA TGTTCCGATG TCGATCACAG GATTGTGATT GTTAGTCCAC TGTATCTCTG 420
ATCGAATAGG TCTATATATT ATTATTTAGA TAGAAAAAGT AGCAATCCAC TTAGGAGATT 480
TATTGATCTG C 491

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 492 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

GCAGATCAAT AAATCTCCTA AGTGGATTGC TACTTTTCT ATCTAAATAA TAATATATAG      60
ACCTATTCGA TCAGAGATAC AGTGGACTAA CAATCACAAT CCTGTGATCG ACATCCGAAC      120
ATAAGTCACT ATCTATCAGA ATAAACAATG ATCCAACGAA TAATAGAGGA GTAAGTGGAC      180
ATGTCCAAAG CATCAGGTAT CATCATGATC GAGAACCACT GTCAAGCAAG ACACAAACAA      240
ACAAAACAGC TTTACACACA AGTCAGCAGT CCAAGCGTTC ATGTCCCAAG GTCAAATGTA      300
ATCATCTGT CATAAGTCCT CGGTCATTGA AAACCAATA AATCGTCGTT TAAACGCGCT      360
CCACGAATTG AGCATTTCCT GTATTCTTCC TTGTACTGGG GCTCTTCATG CTATCCAAGC      420
CAAGACTCTT GCCACTCGTC CCGGTGAACG TTCTCGCAAT GGTGAAGAAG CCCTGTCTTA      480
AACCGAGTTC AT

```

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 492 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

ATGAACTCGG TTTAAGACAG GGCTTCTTCA CCATTGCGAG AACGTTACC GGGACGAGTG      60
GCAAGAGTCT TGGCTTGGAT AGCATGAAGA GCCCCAGTAC AAGGAAGAAT ACTGGAAATG      120
CTCAATTCGT GGAGCGCGTT TAAACGACGA TTTATTTGGT TTTCAATGAC CGAGGACTTA      180
TGACAGGATG ATTACATTG ACCTTGGGAC ATGAACGCTT GGACTGCTGA CTTGTGTGTA      240
AAGCTGTTTT GTTTGTTTGT GTCTTGCTTG ACAGTGGTTC TCGATCATGA TGATACCTGA      300
TGCTTTGGAC ATGTCCACTT ACTCCTCTAT TATTCGTTGG ATCATTGTTT ATTCTGATAG      360
ATAGTGACTT ATGTTCCGAT GTCGATCACA GGGTTGTGAT TGTTAGTCCA CTGTATCTCT      420
GATCGAATAG GTCTATATAT TATTATTTAG ATAGAAAAG TAGCAATCCA CTTAGGAGAT      480
TTATTGATCT GC

```

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 492 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

ATGAACTCGG TTTAAGACAG GGCTTCTTCA CCATTGCGAG AACGTTACC GGGACGAGTG      60
GCAAGAGTCT TGGCTTGGAT AGCATGAAGA GCCCCAGTAC AAGGAAGAAT ACTGGAAATG      120
CTCAATTCGT GGAGCGCGTT TAAACGACGA TTTATTTGGT TTTCAATGAC CGAGGACTTA      180
TGACAGGATG ATTACATTG ACCTTGGGAC ATGAACGCTT GGACTGCTGA CTTGTGTGTA      240
AAGCTGTTTT GTTTGTTTGT GTCTTGCTTG ACAGTGGTTC TCGATCATGA TGATACCTGA      300
TGCTTTGGAC ATGTCCACTT ACTCCTCTAT TATTCGTTGG ATCATTGTTT ATTCTGATAG      360
ATAGTGACTT ATGTTCCGAT GTCGATCACA GGATTGTGAT TGTTAGTCCA CTGTATCTCT      420
GATCGAATAG GTCTATATAT TATTATTTAG ATAGAAAAG TAGCAATCCA CTTAGGAGAT      480
TTATTGATCT GC

```

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 492 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```

ATGAAC TCGG TTTAAG ACAG GGCTTC TTCA CCATTG CCGAG AACGTT CACC GGGACG AGTG      60
GCAAGAG TCT TGGCTT GGAT AGCATG AAGA GCCCCAG TAC AAGGAAGA AT ACTGGAG ATG      120
CTCAATT CGT GGAGCG CGTT TAAACG ACGA TTTATT TGGT TTTCAAT GAC CGAGGAC TTA      180
TGACAGG ATG ATTACATT TG ACCTTG GGAC ATGAAC GCTT GGACTG CTGA CTTGTG TGTA      240
AAGCTGT TTT GTTTGTTT GT GTCTTG CTTG ACAGTG GTTC TCGATC ATGA TGATAC CTGA      300
TGCTTTG GAC ATGTCC ACTT ACTCCC TAT TATTCG TTGG ATCATT GTTT ATTCTG ATAG      360
ATAGTG ACTT ATGTTT CGGAT GTCGAT CACA GGATTG TGAT TGTTAG TCCA CTGTAT CTCT      420
GATCGA ATAG GTCTAT ATAT TATTATT TAG ATAGAAA AG TAGCAAT CCA CTTAGG AGAT      480
TTATTG ATCT GC                                         492

```

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 826 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

TTTAGG CTTG TGCAGG GGAC TCTGTAC ATG TGC GTTGGCA TTATGG ATCG ATTTTAC AG      60
GTTCAG CCAG TTTCCG GAA GAAGCTT CAA TTAGTT GGGG TACTGCT CT GCTCTT GGCC      120
TCCAAG TATG AGGAGAT GTT TTCTCCA AAT ATTGAAG ACT TTGTTT ACAT CACAGACA AT      180
GCTTATA CCA GTTCCCA AAT CCGAGAA ATG GAAACT CTA TTTGAA AGA ATCGAA ATTT      240
GAGTTGG GTC GACCC TTGCC ACTACACT TC TTAAGG CGAG CATCAA AAGC CGGGGAG GTT      300
GATGTTG AAC AGCACACT TT AGCCAAG TAT TTGATG GGAGC TGACTCT CAT CGACTAT GAT      360
ATGGTGC ATT ATCATC CTTT TAAGGTAG CA GCAGCTG CTT CCTGCTT GTC TCAGAAG GTT      420
CTAGGACA AG GAAAAT GGAA CTAAAGC AG CAGTATT ACA CAGGATA CAC AGAGAAT GAA      480
GTATTGGA AG TCATGCAG CA CATGGCCA AG AATGTGG TGA AAGTAA ATGA AACTTAA CT      540
AAATTCA TCG CCATCAAG AA TAAGTAT GCA AGCAGCAA AC TCCTGA AGAT CAGCATG ATC      600
CCTCAG CTGA ACTCAA AAGC CGTCAA AGAC CTTGCCT CTC CACTGAT AGG AAGGTC CTAG      660
GCTGCC GTG CCCCTG GGGG TGTGTG CTTT ATTGTG ECCT TTTTCT TATT GGTTTAG AAC      720
TCTTGAT TTT GTACATA GTC CTCTGG TCTA TCTCATG AAA CCTCTT CTCA GACCA GTTTT      780
CTAAACA TAT ATTGAGG AAA AATAAAG CGA TTGGTT TTTT TTAAGG                                         826

```

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 819 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CTGGAGAGAA	AACCCATAAA	TGCCCCGAAT	GTGGGAGAGC	CTTTTTTTAT	CAGTCATTCC	60
TTATGAGACA	TATGAAATT	CACACTGGAG	AGAAACCGTA	TGAATGTGGG	AAATGTGGGA	120
AAGCCTTTAG	ATATTCCTTA	CACCTTAATA	AACATTTAAG	AAAGCATGTT	GTGCAGAAGA	180
AGCCCTACGA	ATGTGAAGAA	TGTGGGAAAG	TCATTCCGGA	GTCCTCAAAA	TATACACATA	240
TAAGGAGCCA	CACTGGAGAG	AAACCCTATA	AATGTAAGAC	ATGTGGAAAA	GACTTTGCAA	300
AGTCGCCAGG	ACTTAAAAAA	CATCTTAAGA	CTCACAAAGA	TGAGAAGCCC	TGTGAATGAA	360
AGGAAGGTGG	AAAATTTTTC	ATTAATTTTC	TGACTGTACC	AAACATGTGA	GGAGGACATA	420
TTGGAAGGGA	GCTCAAGGGG	TTAGCATGAG	TGAGAACATC	TTCCCTGAAC	TCTCGTATCT	480
TACAGAAATG	TGAAAAAAA	CCCTGTGAAG	GTAAGTCTA	CAGAAAGCCT	TTCATCTTCA	540
TTCATCTTGA	GTAGACATTT	GTTCTCACCC	TGGAGAGAAA	CTGCGAATCT	AAAAGGAATA	600
TGACAAAGCC	TTCAGCGTGG	TCTCAAATTC	ATGGTTCATA	CAAGAACTCA	CACTGCAGAG	660
ACTCCTTACG	GAAATAAAAA	ATGTAGGAAA	GACCTGCCGG	CCGCGGTGGC	TCATGCCTGT	720
AATCCCAGCA	CTTTGGGAGG	CCGAGGCGGG	CGGATCACGA	GGTCAGGAGA	TCAAGACCAT	780
CCTGGCTAAC	ACGGTGATAC	CCCGTCTCTA	CTAAAAATA			819

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1386 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GTAGATGTTG	TATTTCTACT	TTACAGGTAG	GAAAATGGAG	GCTAAGAAAA	GTTAATTTGT	60
CCGAGGGCCC	TCTGATGATA	GTGAACTGG	GATGGAACCT	CTGCCTGCTT	GCTTCTGAGG	120
TCTGGGCTCC	TAATACTGCT	TCTACTGCCT	CGAGCCAAGA	GATCTACGCC	CTATTAAGCA	180
ATTTGTTGTT	CGATAAATTG	GAAGACACAG	CAGATAAGCA	AACAACCTCA	GCAACCAAGT	240
CAGTTCCTGG	AGTTTCTGAA	TTGTTGGGAC	CAAGGGGCCG	TGCAGAGGTA	ACCACAGCTG	300
GCGTAGTGTG	GTTGAGGTAG	CCCTATTAGC	CTTTTAGTTG	CTGTTACTAA	TTTATTTCTC	360
AGTGGTCAAT	GAACCAATTG	GCCATCAATC	AGCTTTGTGT	ATAGGTCATG	TTCCCATGGC	420
TCTGACCCAG	GTTGCTGCTC	AGAGTTGGCA	TCGTGGCTAA	AATATTACTA	GAGGTCAAAG	480
ATATGTGTGT	GTTTGTGGTT	GATTTAGTCG	AGTGATCTAG	AGGAATCTGA	ACCTTAGAGA	540
CTGAAGAAGA	ACCAGCATT	CTGGGCAATA	ATACTTGAGT	TAAGGAGAGT	GTAGCAAAAC	600
TCTAGGTTAG	CATTGGCAGT	CCCTAGAATT	CAGACTGTAG	GCCTAAATGA	CCCTCAGTCC	660
AGAGCTGTAC	CTAATGAGGA	CAATACATTT	TAATGTGAGT	CCATTCTTAA	CAGCAAAATT	720
TCCTCTTTGC	TTGTCACCAG	GGAAAAATGG	GTTTGCATAG	AAAAGGTGGA	GATTGAGGGG	780
GAAGCAGAAT	GGACAAGGAG	TAAAGAGGGA	ATCCAACCTAC	TTAGATTGTA	GCTTTCGTTT	840
TTCTTTGGTA	GTTGTAGAGG	TGAGCTTACC	AAAGCATAGA	TGACAGGCAA	TGTGGTATAC	900
AAGTTACTAC	ACTCCAAAAG	TCTGGGGTTC	TTACTTATTT	TGTGCATGAC	ATCCAAAGTA	960
GCCTAATAAA	ATCTTTTCAC	AGAAAAAAA	GCTTTACTTT	CCTTTGCCAA	ATTTTAACT	1020
TTTTATTCTG	AAATAATTTT	AGAATTATTG	AAAAATTTAG	AGACTAGGAC	AACCCAGATT	1080
CCTCAAATAT	TAACACTTTA	CCACATCTGC	CTTCTCATTC	CTCTCTATAT	ACATAGGTGC	1140
ATGTGTGGTT	TTAATGTTTA	TTTATATACA	TATCATTATT	ATTTTCTTAA	CTGTTTGAGA	1200
GTAAGTTGAA	GACATGATGC	TCCTTACTCT	TTAAATACTT	CAGTGTGTAT	TTCTTAAAAA	1260
GCAGGCCATG	TTCTACATCA	TCACAGTATA	ATTATCAAAA	TTGGGAAATT	AATATTAATG	1320
CAATACTATT	TATCAAATTT	TAAGATCTTA	TTCAAATTTT	ACTTGCTGGC	CTAATAATGT	1380
TCTTTC						1386

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 547 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AAAAGTCTGC	TTTGAGGCAA	AGGTAACCCA	GAATCTCCCA	ATGAAAGAAG	GCTGCACAGA	60
GGTCTCTCTC	CTTCGAGTTG	GGTGGTCTGT	TGATTTTTC	CGTCCACAGC	TTGGTGAAGA	120
TGAATTCTCT	TACGGTTTCG	ATGGACGAGG	ACTCAAGGCA	GAAAATGGAC	AATTTGAGGA	180
ATTTGGCCAG	ACTTTTGGGG	AGAATGATGT	TATTGGCTGC	TTTGCTAATT	TTGAGACTGA	240
AGAAGTAGAA	CTTTCCTTCT	CCAAGAATGG	AGAAGACCTA	GGTGTGGCAT	TCTGGATCAG	300
CAAGGATTCC	CTGGCAGACC	GGGCCCTTCT	ACCCCATGTC	CTCTGCAAAA	ATTGTGTTGT	360
AGAATTAAAC	TTTCGGTCAGA	AGGAGGAGCC	CTTCTTCCCA	CCACCAGAAG	AGTTTGTGTT	420
CATTGATGCT	GTGCCTGTTG	AGGAGCGTGT	ACGCACTGCA	GTCCCTCCCA	AGACCACAGA	480
GGAATGTGAG	GTGATTCTGA	TGGTGGGACT	ACCCGGATCT	GGAAAGACCC	AGTGGGCACT	540
GAAATAT						547



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47, 16/18, G01N 33/53, C12Q 1/68, C12N 15/11, 9/00	A3	(11) International Publication Number: WO 99/67386 (43) International Publication Date: 29 December 1999 (29.12.99)
(21) International Application Number: PCT/US99/14036 (22) International Filing Date: 22 June 1999 (22.06.99) (30) Priority Data: 60/090,391 23 June 1998 (23.06.98) US 60/118,570 3 February 1999 (03.02.99) US 09/337,171 21 June 1999 (21.06.99) US (71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US). (72) Inventor: KENNEDY, Giulia, C.; 360 Castenada Avenue, San Francisco, CA 94116 (US). (74) Agents: POTTER, Jane, E., R.; Chiron Corporation, 4560 Horton Street, Emeryville, CA 94608 (US) et al.	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 15 June 2000 (15.06.00)	
(54) Title: DIFFERENTIALLY EXPRESSED GENES IN PANCREATIC CANCER AND DYSPLASIA (57) Abstract The present invention provides the art with the DNA coding sequences of polynucleotides that are up- or down-regulated in cancer and dysplasia. These polynucleotides and encoded proteins or polypeptides can be used in the diagnosis or identification of cancer and dysplasia. Inhibitors of the up-regulated polynucleotides and proteins can decrease the abnormality of cancer and dysplasia. Enhancing the expression of down-regulated polynucleotides or introducing down-regulated proteins to cells can decrease the growth and/or abnormal characteristics of cancer and dysplasia.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No.

PC, JS 99/14036

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/12 C07K14/47 C07K16/18 G01N33/53 C12Q1/68
C12N15/11 C12N9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K G01N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. AC004934, 15 June 1998 (1998-06-15) LAMAR, B. ET AL.: "Homo sapiens PAC clone DJ0953B05 from 7p12-p14, complete sequence." XP002126839 see nt. 125951-125969 ---	1,2
X	DATABASE EMBL - EMEST3 [Online] Entry/Acc.no. AA775961, 6 February 1998 (1998-02-06) HILLIER, L. ET AL.: "ae78c05.s1 Stratagene schizo brain S11 Homo sapiens cDNA clone 970280 3'." XP002126840 the whole document --- -/--	1,2

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

11 January 2000

Date of mailing of the international search report

17. 04. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Smalt, R

INTERNATIONAL SEARCH REPORT

International Application No

PC. JS 99/14036

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. 004842, 15 June 1998 (1998-06-15) WATERSTON, R.H.: "Homo sapiens clone DJ0609N19, complete sequence." XP002126841 see nt. 111252-111264	1,2
X	DATABASE EMBL EMHUM1 [Online] Entry/acc.no. AC002509, 29 August 1997 (1997-08-29) BIRREN, B. ET AL.: "Homo sapiens chromosome Y, clone 2Y, complete sequence." XP002126842 see nt.99287-99312.	1
X	DATABASE EMBL- EMHUM1 [Online] Entry/Acc.no. AC004168, 24 February 1998 (1998-02-24) STONE, N. E. ET AL.: "Homo sapiens chromosome 4 clone B159L21 map 4q25, complete sequence." XP002126843 the whole document	1
X	WO 98 02560 A (BIOLOG MOLECULAIRE DES PLANTES ;BROUGHTON WILLIAM JOHN (CH); PERRE) 22 January 1998 (1998-01-22) see nt. 136894-136910 of seq.ID.1	1
X	DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. AC004636, 5 May 1998 (1998-05-05) KIMMERLY, W. ET AL.: "Homo sapiens chromosome 5, P1 clone 1268h6 (LBNL H50), complete sequence." XP002126844 see nt. 20805-20821	1
X	DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. AC002488, 25 August 1997 (1997-08-25) BIRREN, B ET AL.: "Homo sapiens chromosome X, clone 36X, complete sequence." XP002126845 see nt. 56517-56531	1

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PL., US 99/14036

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL - EMHUM2. [Online] Entry HS24018, Acc.no. AL021808, 8 February 1998 (1998-02-08) PHILLIPS, S.: "Human DNA sequence from clone 24018 on chromosome 6p21.31-22.2 Contains zinc finger protein pseudogene, VNO-type olfactory receptor pseudogene, nuclear envelope pore membrane protein, EST, STS, GSS." XP002126846 see nt. 63210-63382</p>	1
X	<p>DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. AC004673, 11 May 1998 (1998-05-11) MUZY, D. ET AL.: "Homo sapiens Xp22-132-134 BAC GSHB-590J15 (Genome Systems Human BAC library) complete sequence." XP002126847 see nt. 28491-28700</p>	1
A	<p>EP 0 376 746 A (WISTAR INST) 4 July 1990 (1990-07-04) abstract</p>	
A	<p>WO 86 02081 A (CENTRE NAT RECH SCIENT) 10 April 1986 (1986-04-10) abstract</p>	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/14036

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority; namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Claims 4,6,14,17 and 1-3,5,7,16,19-21 partially.

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 4,6,14,17 and 1-3,5,7,13,16,19-21 partially

Nucleic acids encoding genes which are differentially regulated in pancreatic dysplasia as represented by the seq.ID's 1-11 and 13-15, proteins encoded thereby, antibodies against said proteins, nucleic acid probes derived from said nucleic acids, method for diagnosing dysplasia using said nucleic acids or said proteins, therapeutic compositions comprising said nucleic acids, their complements, said proteins, antibodies thereto, or a ribozyme which binds said nucleic acids.

2. Claims: 8-12,15,18 and 1-3,5,7,13,16,19-21 partially

Nucleic acids encoding a gene which is differentially regulated in pancreatic cancer as represented by the seq.ID 12, the protein encoded thereby, antibodies against said protein, nucleic acid probes derived from said nucleic acid, method for diagnosing cancer using said nucleic acid or said protein, therapeutic compositions comprising said nucleic acid, a complement thereof, said protein, antibodies thereto, or a ribozyme which binds said nucleic acid.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC, JS 99/14036

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9802560 A	22-01-1998	EP 0818465 A	14-01-1998
		EP 0917582 A	26-05-1999
EP 0376746 A	04-07-1990	US 5185254 A	09-02-1993
		AU 4733989 A	16-08-1990
		CA 2006812 A	29-06-1990
		JP 3080081 A	04-04-1991
WO 8602081 A	10-04-1986	FR 2571146 A	04-04-1986
		AT 43610 T	15-06-1989
		EP 0180496 A	07-05-1986
		EP 0197974 A	22-10-1986
		JP 62500304 T	05-02-1987
		OA 8334 A	29-02-1988
		US 4843019 A	27-06-1989



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 15/12, C07K 14/47, 16/18, G01N 33/53, C12Q 1/68, C12N 15/11, 9/00		(11) International Publication Number: WO 99/67386
A3		(43) International Publication Date: 29 December 1999 (29.12.99)
(21) International Application Number: PCT/US99/14036		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 22 June 1999 (22.06.99)		
(30) Priority Data:		
60/090,391	23 June 1998 (23.06.98) US	
60/118,570	3 February 1999 (03.02.99) US	
09/337,171	21 June 1999 (21.06.99) US	
(71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US).		Published <i>With a revised version of the international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(72) Inventor: KENNEDY, Giulia, C.; 360 Castenada Avenue, San Francisco, CA 94116 (US).		
(74) Agents: POTTER, Jane, E., R.; Chiron Corporation, 4560 Horton Street, Emeryville, CA 94608 (US) et al.		
		(88) Date of publication of the international search report: 15 June 2000 (15.06.00)
		(88) Date of publication of the revised version of the international search report: 20 July 2000 (20.07.00)
(54) Title: DIFFERENTIALLY EXPRESSED GENES IN PANCREATIC CANCER AND DISPLASIA		
(57) Abstract		
<p>The present invention provides the art with the DNA coding sequences of polynucleotides that are up- or down-regulated in cancer and dysplasia. These polynucleotides and encoded proteins or polypeptides can be used in the diagnosis or identification of cancer and dysplasia. Inhibitors of the up-regulated polynucleotides and proteins can decrease the abnormality of cancer and dysplasia. Enhancing the expression of down-regulated polynucleotides or introducing down-regulated proteins to cells can decrease the growth and/or abnormal characteristics of cancer and dysplasia.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Intern: al Application No
PCT/US 99/14036

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/12 C07K14/47 C07K16/18 G01N33/53 C12Q1/68
C12N15/11 C12N9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N C07K G01N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. AC004934, 15 June 1998 (1998-06-15) LAMAR, B. ET AL.: "Homo sapiens PAC clone DJ0953B05 from 7p12-p14, complete sequence." XP002126839 see nt. 125951-125969	1,2
X	DATABASE EMBL - EMEST3 [Online] Entry/Acc.no. AA775961, 6 February 1998 (1998-02-06) HILLIER, L. ET AL.: "ae78c05.s1 Stratagene schizo brain S11 Homo sapiens cDNA clone 970280 3'." XP002126840 the whole document	1,2

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

25 April 2000

Date of mailing of the international search report

22.05.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Smalt, R

INTERNATIONAL SEARCH REPORT

Intern. Application No.

PCT/US 99/14036

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages.	Relevant to claim No.
X	<p>DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. 004842, 15 June 1998 (1998-06-15) WATERSTON, R.H.: "Homo sapiens clone DJ0609N19, complete sequence." XP002126841 see nt. 111252-111264</p>	1,2
X	<p>DATABASE EMBL EMHUM1 [Online] Entry/acc.no. AC002509, 29 August 1997 (1997-08-29) BIRREN, B. ET AL.: "Homo sapiens chromosome Y, clone 2Y, complete sequence." XP002126842 see nt.99287-99312.</p>	1
X	<p>DATABASE EMBL- EMHUM1 [Online] Entry/Acc.no. AC004168, 24 February 1998 (1998-02-24) STONE, N. E. ET AL.: "Homo sapiens chromosome 4 clone B159L21 map 4q25, complete sequence." XP002126843 the whole document</p>	1
X	<p>WO 98 02560 A (BIOLOG MOLECULAIRE DES PLANTES ;BROUGHTON WILLIAM JOHN (CH); PERRE) 22 January 1998 (1998-01-22) see nt. 136894-136910 of seq.ID.1</p>	1
X	<p>DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. AC004636, 5 May 1998 (1998-05-05) KIMMERLY, W. ET AL.: "Homo sapiens chromosome 5, P1 clone 1268h6 (LBNL H50), complete sequence." XP002126844 see nt. 20805-20821</p>	1
X	<p>DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. AC002488, 25 August 1997 (1997-08-25) BIRREN, B ET AL.: "Homo sapiens chromosome X, clone 36X, complete sequence." XP002126845 see nt. 56517-56531</p>	1

-/--

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/US 99/14036

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL - EMBEST3 [Online] Entry/Acc.no. AA812053, 16 February 1998 (1998-02-16) STRAUSBERG, R.: "ob48c07.s1 NCI CGAP GC81 Homo sapiens cDNA clone IMAGE:1334604.3 similar to SW:CGB2 MESAU P37883 G2/MITOTIC-SPECIFIC CYCLIN B2. ;," XP002136369 the whole document	1
X	DATABASE EMBL - EMHUM2 [Online] Entry HS24018, Acc.no. AL021808, 8 February 1998 (1998-02-08) PHILLIPS, S.: "Human DNA sequence from clone 24018 on chromosome 6p21.31-22.2 Contains zinc finger protein pseudogene, VNO-type olfactory receptor pseudogene, nuclear envelope pore membrane protein, EST, STS, GSS." XP002126846 see nt. 63210-63382	1
X	DATABASE EMBL - EMHUM1 [Online] Entry/Acc.no. AC004673, 11 May 1998 (1998-05-11) MUZNY, D. ET AL.: "Homo sapiens Xp22-132-134 BAC GSHB-590J15 (Genome Systems Human BAC library) complete sequence." XP002126847 see nt. 28491-28700	1
A	EP 0 376 746 A (WISTAR INST) 4 July 1990 (1990-07-04) abstract	
A	WO 86 02081 A (CENTRE NAT RECH SCIENT) 10 April 1986 (1986-04-10) abstract	
P,X	WO 98 55510 A (COCKS BENJAMIN GRAEME ; INCYTE PHARMA INC (US); CORLEY NEIL C (US);) 10 December 1998 (1998-12-10) whole document, particularly the claims relating to HCRP-2.	1,3,5

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/14036

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 4,6,14,17 and 1-3,5,7,13,16,19-21 partially

Nucleic acids encoding genes which are differentially regulated in pancreatic dysplasia as represented by the seq.ID's 1-11 and 13-15, proteins encoded thereby, antibodies against said proteins, nucleic acid probes derived from said nucleic acids, method for diagnosing dysplasia using said nucleic acids or said proteins, therapeutic compositions comprising said nucleic acids, their complements, said proteins, antibodies thereto, or a ribozyme which binds said nucleic acids.

2. Claims: 8-12,15,18 and 1-3,5,7,13,16,19-21 partially

Nucleic acids encoding a gene which is differentially regulated in pancreatic cancer as represented by the seq.ID 12, the protein encoded thereby, antibodies against said protein, nucleic acid probes derived from said nucleic acid, method for diagnosing cancer using said nucleic acid or said protein, therapeutic compositions comprising said nucleic acid, a complement thereof, said protein, antibodies thereto, or a ribozyme which binds said nucleic acid.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/14036

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9802560 A	22-01-1998	EP 0818465 A	14-01-1998
		EP 0917582 A	26-05-1999
EP 0376746 A	04-07-1990	US 5185254 A	09-02-1993
		AU 4733989 A	16-08-1990
		CA 2006812 A	29-06-1990
		JP 3080081 A	04-04-1991
WO 8602081 A	10-04-1986	FR 2571146 A	04-04-1986
		AT 43610 T	15-06-1989
		EP 0180496 A	07-05-1986
		EP 0197974 A	22-10-1986
		JP 62500304 T	05-02-1987
		OA 8334 A	29-02-1988
		US 4843019 A	27-06-1989
WO 9855510 A	10-12-1998	AU 7954998 A	21-12-1998